



## Nutrition Spotlight: U.S. Dairy Proteins: High-quality and Complete Protein Sources



All dairy foods and ingredients start out as milk, which contains 3.5 percent protein — 80 percent casein and 20 percent whey. Milk protein ingredients contain the same protein ratio found in milk, while whey protein ingredients contain 100 percent whey. U.S. dairy proteins with protein levels 80 percent and higher are widely used in health and wellness and sports nutrition products.

### Benefits of Dairy Protein

Published nutrition research on the health and nutritional benefits of dairy proteins continues to rise each year, supporting the benefits of incorporating whey and milk proteins into the daily diet. Protein is an essential nutrient the body needs to build and maintain muscle. Protein also plays an integral role in the body's structure, functions and regulation of all tissues and organs.

There are many health and wellness benefits for all types of lifestyles that come from eating U.S. dairy proteins. Research shows that<sup>1-18</sup> diets higher in protein help slow muscle loss, curb hunger, maintain a healthy weight, build lean muscle (with regular resistance exercise) and enhance exercise recovery.

#### DID YOU KNOW

- U.S. dairy proteins score at or near the top in current methods used to measure protein quality.
- The United Nations Food and Agriculture Organization (FAO) has recommended a new method to assess protein quality by its ability to supply the body with amino acids.
- U.S. whey protein is high in branched-chain amino acids and leucine content, making it unique in its ability to initiate new muscle synthesis.



Whether aiming to increase lean muscle or add muscle mass, eating U.S. dairy proteins before or after exercise can help improve body composition.



U.S. dairy proteins can help people maintain muscle as they age.



Weekend warriors, sports enthusiasts and hard-core athletes can meet their goals and recover faster from exercise with help from U.S. dairy proteins.




U.S. dairy proteins can help people control their hunger and maintain a healthy weight.

## Not All Proteins Are Created Equal

High-quality protein — such as U.S. dairy protein — contains the essential (sourced solely from food) and non-essential (synthesized by the body) amino acids the body needs. In contrast, many plant-based proteins used as ingredients are not considered complete sources as they do not contain all of the essential amino acids.

Studies show that essential amino acids can maximally stimulate muscle protein synthesis at relatively low amounts:





## SPOTLIGHT

### Amino Acids

Specific essential amino acids, called branched-chain amino acids (BCAAs), play an even greater role in muscle protein metabolism as, unlike other types of amino acids, these bypass the liver and go straight to the muscle. The result is a faster rate of absorption for recovery, maintenance and growth.<sup>23,24,25</sup> And, among the three BCAAs (leucine, isoleucine, valine), research shows leucine is unique in its ability to initiate new muscle synthesis.<sup>3,26</sup> See how dairy protein sources stack up to other sources:

PROTEIN SOURCE	LEUCINE	BCAA
Whey Protein Isolate	14%	26%
Casein	10%	23%
Milk Protein	10%	21%
Egg Protein	9%	20%
Muscle Protein (Meat, Chicken, etc.)	8%	18%
Soy Protein Isolate	8%	18%
Wheat Protein	7%	15%

Values reflect grams of amino acids/100 grams of protein. Source: USDA Food Composition Tables modified from Layman, DK. The role of leucine in weight loss diets and glucose homeostasis. *J Nutr.* 2003;133(1):261S-267S.



## Measuring Protein Quality

There are a variety of ways in which protein quality is measured. Recently, a new method has been introduced that assesses the quality of dietary proteins by their ability to supply amino acids for use by the body. This method, the Digestible Indispensable Amino Acid Score (DIAAS)<sup>27</sup> is being recommended for use based on a groundbreaking United Nations Food and Agriculture Organization (FAO) Expert Consultation report. While more data is needed to support the full implementation of this new method, this report shows that high-quality proteins from milk, whey and other dairy products may score 30 percent higher compared with scores using older methods. As shown below, dairy proteins already are rated at or near the top of all current measures.

PROTEIN TYPE	PDCAAS*	BIOLOGICAL VALUE	NET PROTEIN UTILIZATION	PROTEIN EFFICIENCY RATIO
Whey Protein	1.00	104	92	3.2
Milk	1.00	91	82	2.5
Casein	1.00	77	76	2.5
Egg	1.00	100	94	3.9
Soy Protein	1.00	74	61	2.2
Beef	0.92	80	73	2.9
Black Beans	0.75		0	0
Peanuts	0.52			1.8
Wheat Gluten	0.25	64	92	0.8

\*Protein digestibility-corrected amino acid score

And, when compared with plant-based proteins, research shows whey protein is more beneficial for muscle growth due to its content of BCAAs, specifically leucine.<sup>28</sup> BCAAs — leucine, isoleucine, valine — are especially beneficial post-workout because, unlike other amino acids, BCAAs bypass the liver and go straight to the muscle tissue, providing a faster rate of absorption for recovery, maintenance and growth.<sup>29,30,28</sup>








## THE RESEARCH

### Dairy Protein vs. Other Sources

Whey protein isolate, casein and milk protein all contain high amounts of BCAAs, and whey protein is one of the richest sources of leucine,<sup>31,32</sup> making it an excellent partner when focusing on weight management. Research shows that the leucine content of protein may be the key to promoting greater gains in lean body mass with resistance exercise.<sup>24</sup> Even without exercise, whey protein has been shown to support increased weight and fat loss.<sup>7</sup>

### A Closer Look at the Science

In a study published in the *Journal of the American College of Nutrition*, 36 men and 27 women completed a nine-month resistance exercise study consuming whey, soy or carbohydrate. An equivalent amount was consumed each day — after their exercise session on training days or at breakfast on nontraining days.



Source: Volek JS, Volk BM, Gomez AL, et al. Whey protein supplementation during resistance training augments lean body mass. *J Am Coll Nutr.* 2013;32(2):122-136.


#### RESULTS

The whey group showed significant increases in lean mass (3.3 kg ± 1.5 kg) over soy (1.8 kg ± 1.6 kg) and carbohydrates (2.3 kg ± 1.7 kg) at every time point tested. The results also showed that the plasma leucine levels doubled after whey consumption but only showed minimal changes in the soy and carbohydrate groups when consumed directly after exercise.

#### WHEY ADVANTAGE

Despite consuming similar calories and protein during resistance training, daily supplementation with whey was more effective than soy protein in promoting gains in lean body mass, highlighting that the extra leucine content in whey is an important factor in lean body mass responses to resistance training.

In another study published in the *Journal of Nutrition*, 73 overweight and obese adults completed a 23-week clinical trial consuming 56 grams per day of whey protein, soy protein or carbohydrate as a beverage in two even doses with breakfast and dinner.



Source: Baer DJ, Stote KS, Paul DR, Harris GK, Rumpler WV, Clevidence BA. Whey protein but not soy protein supplementation alters body weight and composition in free-living overweight and obese adults. *J Nutr.* 2011;141(8):1489-1494.

**RESULTS**

Those who consumed whey protein had considerably less fat mass (2.3 kg lower) than the carbohydrate group. Waist circumference among the whey protein group decreased by 2.4 cm compared with the soy protein and carbohydrate groups, and body weight was 1.8 kg lower in the whey protein group.

**WHEY ADVANTAGE**

Diets higher in protein — in particular, whey protein — may help in long-term maintenance of body weight and composition among overweight and obese individuals even when calories are not restricted.



**GET IN TOUCH**

**Looking for more information on dairy proteins?**

While the U.S. Dairy Export Council® (USDEC) does not manufacture or sell dairy products, we proudly support the people who do. Search [ThinkUSAdairy.org/Nutrition](http://ThinkUSAdairy.org/Nutrition) for more information on the health and nutritional benefits of U.S. dairy proteins.

- <sup>1</sup> Houston D, Nicklas BJ, Ding J, et al. Dietary protein intake is associated with lean mass change in older, community-dwelling adults: the Health, Aging, and Body Composition (Health ABC) study. *Am J Clin Nutr*. 2008;87(1):150-155.
- <sup>2</sup> Mojtahedi M, Thorpe M, Karampinos D, et al. The effects of a higher protein intake during energy restriction on changes in body composition and physical function in older women. *J Gerontol A Biol Sci Med Sci*. 2011;66(11):1218-1225.
- <sup>3</sup> Institute of Medicine. *Macronutrients and healthful diets. Dietary Reference Intakes for Energy, Carbohydrate, Fiber, Fat, Fatty Acids, Cholesterol, Protein, and Amino Acids (Macronutrients) (2005)*. Washington, DC: National Academies Press; 2002/2005:769-876.
- <sup>4</sup> Smeets AJ, Soenen S, Luscombe-Marsh ND, Ueland O, Westerterp-Plantenga M. Energy expenditure, satiety, and plasma ghrelin, glucagon-like peptide 1, and peptide tyrosine-tyrosine concentrations following a single high-protein lunch. *J Nutr*. 2008;138(4):698-702.
- <sup>5</sup> Leidy HJ, Armstrong CL, Tang M, Mattes RD, Campbell WW. The influence of higher protein intake and greater eating frequency on appetite control in overweight and obese men. *Obesity*. 2010;18(9):1725-1732.
- <sup>6</sup> Josse AR, Atkinson SA, Tarnopolsky MA, Phillips SM. Increased consumption of dairy foods and protein during diet- and-exercise-induced weight loss promotes fat mass loss and lean muscle gain in overweight and obese premenopausal women. *J Nutr*. 2011;141(19):1626-1634.
- <sup>7</sup> Baer D, Stote KS, Paul D, Harris G, Rumpler W, Clevidence B. Whey protein but not soy protein supplementation alters body weight and composition in free-living overweight and obese adults. *J Nutr*. 2011;141(8):1489-1494.
- <sup>8</sup> Westerterp-Plantenga MS, Nieuwenhuizen A, Tomé D, Soenen S, Westerterp KR. Dietary protein, weight loss, and weight maintenance. *Annu Rev Nutr*. 2009;29:21-41.
- <sup>9</sup> Claessens M, van Baak MA, Monsheimer S, Saris WH. The effect of a low-fat, high-protein or high-carbohydrate ad libitum diet on weight loss maintenance and metabolic risk factors. *Int J Obes*. 2009;33(3):296-304.
- <sup>10</sup> Westerterp-Plantenga MS, Lejeune MP, Nijs I, van Ooijen M, Kovacs EM. High protein intake sustains weight maintenance after body weight loss in humans. *Int J Obes*. 2004;28(1):57-64.
- <sup>11</sup> Lejeune MP, Kovacs EM, Westerterp-Plantenga MS. Additional protein intake limits weight regain after weight loss in humans. *Br J Nutr*. 2005;93(2):281-289.
- <sup>12</sup> Tang JE, Moore DR, Kujbida GW, Tarnopolsky MA, Phillips SM. Ingestion of whey hydrolysate, casein, or soy protein isolate: effects on mixed muscle protein synthesis at rest and following resistance exercise in young men. *J Appl Physiol*. 2009;107(3):987-992.
- <sup>13</sup> Tang JE, Phillips SM. Maximizing muscle protein anabolism: the role of protein quality. *Curr Opin Clin Nutr Metab Care*. 2009;12(1):66-71.
- <sup>14</sup> Churchward-Benne TA, Burd NA, Mitchell CJ, et al. Supplementation of a suboptimal protein dose with leucine or essential amino acids: effects on myofibrillar protein synthesis at rest and following resistance exercise in men. *J Physiol*. 2012;590(Pt 11):2751-2765.
- <sup>15</sup> Tipton KD, Elliott TA, Cree MG, et al. Ingestion of casein and whey proteins result in muscle anabolism after resistance exercise. *Med Sci Sports Exerc*. 2004;36(12):2073-2081.
- <sup>16</sup> Howarth KR, Moreau NA, Phillips SM, Gibala MJ. Coingestion of protein with carbohydrate during recovery from endurance exercise stimulates skeletal muscle protein synthesis in humans. *J Appl Physiol*. 2009;106(4):1394-1402.
- <sup>17</sup> Tang JE, Manolagos JJ, Kujbida GW, et al. Minimal whey protein with carbohydrate stimulates muscle protein synthesis following resistance exercise in trained young men. *Appl Physiol Nutr Metab*. 2007;32(6):1132-1138.
- <sup>18</sup> Tipton KD, Elliott TA, Cree MG, et al. Stimulation of net muscle protein synthesis by whey protein ingestion before and after exercise. *Am J Physiol Endocrinol Metab*. 2007;292(1):E71-E76.
- <sup>19</sup> Moore DR, Robinson MJ, Fry JL, et al. Ingested protein dose response of muscle and albumin protein synthesis after resistance exercise in young men. *Am J Clin Nutr*. 2009;89(1):1-8.
- <sup>20</sup> Witard OC, Jackman SR, Breen L, et al. Myofibrillar muscle protein synthesis rates subsequent to a meal in response to increasing doses of whey protein at rest and after resistance exercise. *Am J Clin Nutr*. 2014;99(1):86-95.
- <sup>21</sup> Paddon-Jones D, Rasmussen BB. Review: Dietary protein recommendations and the prevention of sarcopenia. *Curr Opin Clin Nutr Metab Care*. 2009;12(1):86-90.
- <sup>22</sup> Pennings B, Groen B, de Lange A, et al. Amino acid absorption and subsequent muscle protein accretion following graded intakes of whey protein in elderly men. *Am J Physiol Endocrinol Metab*. 2012;302(8):E992-E999.
- <sup>23</sup> Greiwe JS, Kwon G, McDaniel ML, Semenkovich CF. Leucine and insulin activate p70 S6 kinase through different pathways in human skeletal muscle. *Am J Physiol Endocrinol Metab*. 2001;281(3):E466-E471.
- <sup>24</sup> Rieu I, Balage M, Sornet C, et al. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol*. 2006;575(Pt 1):305-315.
- <sup>25</sup> Layman DK. The role of leucine in weight loss diets and glucose homeostasis. *J Nutr*. 2003;133(1):261S-267S.
- <sup>26</sup> Pasiakos SM, McClung HL, McClung JP, et al. Leucine-enriched essential amino acid supplementation during moderate steady state exercise enhances postexercise muscle protein synthesis. *Am J Clin Nutr*. 2011;94(3):809-818.
- <sup>27</sup> Food and Agriculture Organization of the United Nations. *Dietary protein quality evaluation in human nutrition*; 2013.
- <sup>28</sup> Volek JS, Volk BM, Gomez AL, et al. Whey protein supplementation during resistance training augments lean body mass. *J Am Coll Nutr*. 2013;32(2):122-136.
- <sup>29</sup> Greiwe JS, Kwon G, McDaniel ML, et al. Leucine and insulin activate p70 S6 kinase through different pathways in human skeletal muscle. *Am J Physiol Endocrinol Metab*. 2001;281(3):E466-E471.
- <sup>30</sup> Rieu I, Balage M, Sornet C, et al. Leucine supplementation improves muscle protein synthesis in elderly men independently of hyperaminoacidaemia. *J Physiol*. 2006;575(Pt1):305-315.
- <sup>31</sup> Layman DK. The role of leucine in weight loss diets and glucose homeostasis. *J Nutr*. 2003;133(1):261S-267S.
- <sup>32</sup> Layman DK, Walker DA. Potential importance of leucine in treatment of obesity and the metabolic syndrome. *J Nutr*. 2006;136(Suppl 1):319S-323S.



To learn more and find a USDEC representative near you, go to [ThinkUSAdairy.org/global-presence](http://ThinkUSAdairy.org/global-presence).